



Non-Native Invasive Plants of the City of Alexandria, Virginia

Non-native invasive plants have increasingly become a major threat to natural areas, parks, forests, and wetlands by displacing native species and wildlife and significantly degrading habitats. Today, they are considered the greatest threat to natural areas and global biodiversity, second only to habitat loss resulting from development and urbanization (Vitousek et al. 1996, Pimentel et al. 2005).

The Virginia Department of Conservation and Recreation has identified 90 non-native invasive plants that threaten natural areas and lands in Virginia (Heffernan et al. 2014) and Swearingen et al. (2010) include 80 plants from a list of nearly 280 non-native invasive plant species documented within the mid-Atlantic region. Largely overlapping with these and other regional lists are 116 species that were documented in the City of Alexandria, Virginia during vegetation surveys and natural resource assessments by the City of Alexandria Dept. of Recreation, Parks, and Cultural Activities (RPCA), Natural Lands Management Section.

This list is not regulatory but serves as an educational reference informing those with concerns about non-native invasive plants in the City of Alexandria and vicinity, including taking action to prevent the further spread of these species by not planting them.

Exotic species are those that are not native to a particular place or habitat as a result of human intervention. A non-native invasive plant is here defined as one that exhibits some degree of invasiveness, whether dominant and widespread in a particular habitat or landscape or much less common but long-lived and extremely persistent in places where it occurs. The presence of non-native invasive plants is largely the result of soil and habitat disturbance, though many of these species are capable of eventually spreading from degraded areas into relatively undisturbed sites, such as interior forest.

Hundreds of exotic plants growing in a variety of habitats occur in the City of Alexandria. Many of these are widespread, early spring-blooming annuals of lawns and open disturbed areas, such as Chickweeds (*Stellaria media*, *Cerastium* spp.), Speedwells (*Veronica* spp.), Purple Dead-nettle (*Lamium purpureum*), Hairy Bittercress (*Cardamine hirsuta*), and others. Some, like Mugwort (*Artemisia vulgaris*), Bermuda Grass (*Cynodon dactylon* var. *dactylon*), and Dandelion (*Taraxacum officinale*), are familiar, difficult to eradicate perennial weeds of lawn and garden. Many others are commonly visible along roadways, medians, cracks in pavement, industrial sites, and other highly disturbed places.

However, only exotic plants that are especially invasive, damaging, and persistent, including species that produce a pervasive seed bank and those that prevent natural succession by native plants, are included on the following list. Native species that are typically weedy in disturbed areas, such as Poison

Ivy (*Toxicodendron radicans*), Black Locust (*Robinia pseudoacacia*), Black Cherry (*Prunus serotina*), Common Pokeweed (*Phytolacca americana*), etc., are not included here because they are not non-native invasive species, they do not degrade natural areas, and they are important for native wildlife.

Non-native invasive plants occurring within the City of Alexandria are actively monitored by RPCA Natural Lands Management Section, which updates the list annually as new data are collected and species are assessed for invasiveness.

Herbaceous Plants

*A species that is native to North America but not to the City of Alexandria, that is now found naturalized in Alexandria outside of its natural range.

Garlic Mustard (*Alliaria petiolata*)
 Joint-head Grass (*Arthraxon hispidus* var. *hispidus*)
 Italian Arum (*Arum italicum*)
 Fountain Grass (*Cenchrus purpurascens*) [= *Pennisetum alopecuroides*]
 Knapweed (*Centaurea* spp.)
 Canada Thistle (*Cirsium arvense*)
 Common Thistle (*Cirsium vulgare*)
 Variable Flatsedge (*Cyperus difformis*)
 Helleborine (*Epipactis helleborine*)
 Weeping Lovegrass (*Eragrostis curvula*)
 Mulberry-weed (*Fatoua villosa*)
 Lesser Celandine (*Ficaria verna*) [= *Ranunculus ficaria*]
 Ground-ivy (*Glechoma hederacea*)
 Orange Day-lily (*Hemerocallis fulva*)
 Hydrilla (*Hydrilla verticillata*)
 Yellow Iris (*Iris pseudacorus*)
 Korean-clover (*Kummerowia stipulacea*)
 Japanese-clover (*Kummerowia striata*)
 Pasture Spikesedge (*Kyllinga gracillima*)
 Yellow Archangel (*Lamium galeobdolon*) [= *Lamiastrum galeobdolon*]
 Chinese Lespedeza (*Lespedeza cuneata*)
 Big Blue Lilyturf (*Liriope muscari*)
 Creeping Lilyturf (*Liriope spicata*)
 Floating Primrose-willow (*Ludwigia peploides* var. *glabrescens*)
 Annual Honesty, Money Plant (*Lunaria annua*)
 Purple Loosestrife (*Lythrum salicaria*)
 Japanese Stiltgrass (*Microstegium vimineum*)
 Chinese Silvergrass, Eulalia (*Miscanthus sinensis*)
 Marsh Dewflower (*Murdannia keisak*) [= *Aneilema keisak*]
 Parrot Feather (*Myriophyllum aquaticum*)
 Eurasian Water-milfoil (*Myriophyllum spicatum*)

Java Dropwort (*Oenanthe javanica*)
 Common Star-of-Bethlehem (*Ornithogalum umbellatum*)
 Dallis Grass (*Paspalum dilatatum*)
 Beefsteak Plant (*Perilla frutescens*)
 Long-bristled Smartweed (*Persicaria longiseta*) [= *Polygonum cespitosum*]
 Common Reed (*Phragmites australis* ssp. *australis*)
 Japanese Knotweed (*Reynoutria japonica*) [= *Polygonum cuspidatum*]
 Giant Knotweed (*Reynoutria sachalinensis*) [= *Polygonum sachalinense*]
 Curly Dock (*Rumex crispus* ssp. *crispus*)
 Crown Vetch (*Securigera varia*) [= *Coronilla varia*]
 Yellow Foxtail (*Setaria pumila* ssp. *pumila*)
 Green Foxtail (*Setaria viridis* var. *viridis*)
 Johnson Grass (*Sorghum halepense*)
 Vetch (*Vicia* spp.)
 Oriental False Hawksbeard (*Youngia japonica*) [= *Crepis japonica*]

Vines

Five-leaf Akebia (*Akebia quinata*)
 Porcelain-berry (*Ampelopsis brevipedunculata*)
 Oriental Bittersweet (*Celastrus orbiculatus*)
 Sweet Autumn Clematis (*Clematis terniflora*)
 Chinese Yam (*Dioscorea polystachya*) [= *Dioscorea batatas*]
 Winter Creeper (*Euonymus fortunei*)
 English Ivy (*Hedera helix*) [Recent morphological and flow cytometry studies of *Hedera* spp. in the greater D.C. area have shown diploid *H. helix* to be by far the most common naturalized *Hedera* species in the region, with rare occurrences of tetraploid *H. hibernica* present (McAllister and Simmons, in prep.). Flow cytometry studies by Alan Whittemore and Jun Wen have also found triploids in areas where *H. helix* and *H. hibernica* grow together (Alan Whittemore, pers. comm.)]
 Japanese Honeysuckle (*Lonicera japonica*)
 Boston-ivy (*Parthenocissus tricuspidata*)
 Mile-a-minute Weed (*Persicaria perfoliata*) [= *Polygonum perfoliatum*]
 Kudzu (*Pueraria montana*)
 Greater Periwinkle (*Vinca major*)
 Common Periwinkle (*Vinca minor*)
 Japanese Wisteria (*Wisteria floribunda*)
 Chinese Wisteria (*Wisteria sinensis*) [“Some of our material may represent *Wisteria* x *formosa* Rehder, a complex series of hybrids and backcrosses between *W. floribunda* and *W. sinensis*.” (VBA 2019)]

Shrubs

Leatherleaf Mahonia (*Berberis bealei*) [= *Mahonia bealei*]
 Japanese Barberry (*Berberis thunbergii*)
 Butterfly-bush (*Buddleia davidii*)

Hardy Orange (*Citrus trifoliata*) [= *Poncirus trifoliata*]
 Thorny Olive (*Elaeagnus pungens*)
 Autumn Olive (*Elaeagnus umbellata*)
 Burning Bush (*Euonymus alatus*)
 Rose-of-Sharon (*Hibiscus syriacus*)
 Chinese Holly (*Ilex cornuta*)
 Japanese Holly (*Ilex crenata*)
 Privet (*Ligustrum* spp.) [Japanese Privet (*Ligustrum japonicum*), Border Privet (*Ligustrum obtusifolium* var. *obtusifolium*), California Privet (*Ligustrum ovalifolium*), Chinese Privet (*Ligustrum sinense*), and Common Privet (*Ligustrum vulgare*) all occur in Alexandria, as well as likely others.]
 Amur honeysuckle (*Lonicera maackii*)
 Nandina (*Nandina domestica*)
 Holly Osmanthus (*Osmanthus heterophyllus*)
 Golden Bamboo (*Phyllostachys aurea*)
 Yellow Groove Bamboo (*Phyllostachys aureosulcata*) [Yellow Groove Bamboo is frequently confused for Golden Bamboo (*Phyllostachys aurea*), when in fact Yellow Groove Bamboo is far more common in northern Virginia and the Washington, D.C. region.]
 Bisset's Bamboo, Hardy Bamboo (*Phyllostachys bissetii*)
 Arrow Bamboo (*Pseudosasa japonica*)
 Multiflora Rose (*Rosa multiflora*)
 Himalayan Blackberry (*Rubus discolor*) [Eedes and Newton (1988) state that *R. ulmifolius* is probably the only variable sexual species in subgenus *Rubus*. *R. bifrons* is given as of dubious attribution, and *R. discolor* and *R. armeniacus* as synonyms of *R. procerus* Mueller ex Boulay, *Ronces Vosg.* 7 (1864), with the comment that it is probably a widespread species native to Europe with 2n=28 (so tetraploid and apomictic), which was probably sent from Germany to the U.S. nurseries where it was given the name 'Himalayan Giant' and exported back to Europe (Hugh McAllister, pers. comm.). According to Eedes and Newton, *R. procerus/discolor* "is readily bird-sown and quickly forms dense patches of very robust plants" and is "usually distinguished from related species without difficulty by its large size."]
 Wineberry (*Rubus phoenicolasius*)
 Linden Arrow-wood (*Viburnum dilatatum*)
 Tea Viburnum (*Viburnum setigerum*)

Trees

Norway Maple (*Acer platanoides*)
 Tree-of-Heaven (*Ailanthus altissima*)
 Mimosa, Silk Tree (*Albizia julibrissin*)
 Golden Rain Tree (*Koelreuteria paniculata*)
 Siberian Crabapple (*Malus baccata*) [Includes the likely many hybrids and backcrosses between *M. baccata* and *M. floribunda* and other exotic *Malus* species]
 Japanese Flowering Crabapple (*Malus floribunda*)
 Plumleaf Crabapple (*Malus prunifolia*)
 Siebold's Crabapple (*Malus sieboldii*)

White Mulberry (*Morus alba*)
 Princess Tree (*Paulownia tomentosa*)
 Sweet Cherry (*Prunus avium*)
 Higan Cherry (*Prunus subhirtella*)
 Callery Pear, Bradford Pear (*Pyrus calleryana*) [Includes all named varieties and cultivars of *P. calleryana*, such as “Aristocrat”, “Autumn Blaze”, “Chanticleer”, “Cleveland Select”, “Redspire”, Whitehouse”, etc.]
 Sawtooth Oak (*Quercus acutissima*)
 White Willow (*Salix alba*)
 Crack Willow (*Salix x fragilis*) [= *Salix x rubens*; ‘Prior to the lectotypification of *Salix fragilis* L. and the description of *S. euxina* (Belyaeva 2009) – formerly *S. fragilis* - the name *S. "fragilis"* was often inadvertently used for both the pure species and for its hybrids with *S. alba*.’ (FNA 2010). The presence of *S. euxina* in Alexandria is unknown.]
 Pagoda Tree (*Styphnolobium japonicum*) [= *Sophora japonica*]
 Siberian Elm (*Ulmus pumila*)

Invasive Watchlist

*Big Bluestem (*Andropogon gerardii*) [Frequently planted in our region as a component of generic native meadow seed mixes, or as an accidental contaminant in mixes. In northeastern Virginia it is localized and native only to riverside prairies and outcrops of the Potomac Gorge in Fairfax County.]
 *Meadow Sedge (*Carex flaccosperma*) [This taxon is not native to anywhere near Alexandria; “in the eastern U.S., it ranges as far north as the southeastern portion of Virginia” (Rob Naczi, pers. comm.). It is commonly available through the nursery industry and is popular for its bluish-green foliage.]
 Japanese Grass Sedge, Morrow’s Sedge (*Carex morrowii*) [Includes the cultivars ‘Variegata’ and ‘Ice Dance’ that are common in the landscape industry.]
 Chinese Chestnut (*Castanea mollissima*)
 Japanese Shield Fern (*Dryopteris erythrosora*)
 European Spindle Tree (*Euonymus europaeus*)
 Castor Aralia (*Kalopanax septemlobus*)
 *Southern Magnolia (*Magnolia grandiflora*)
 Cherry Laurel (*Prunus laurocerasus*)
 Ravenna-grass (*Tripidium ravennae*) [= *Saccharum ravennae*; *Erianthus ravennae*]
 Japanese Snowball (*Viburnum plicatum*)
 Leatherleaf Viburnum (*Viburnum rhytidophyllum*)
 Tall False Hawksbeard (*Youngia thunbergiana*)
 Japanese Zelkova (*Zelkova serrata*)

References

Brewer, J.S. 2010. A potential conflict between preserving regional plant diversity and biotic resistance to an invasive grass, *Microstegium vimineum*. *Natural Areas Journal* 30: 279-293.

- Cooley, G. 1994. Invasive exotic plants that threaten native species and natural habitats in Maryland. Maryland Department of Natural Resources, Natural Heritage Program. Annapolis, Maryland.
- Culley, T.M. and N.A. Hardman. 2007. The beginning of a new invasive plant: A history of the ornamental callery pear in the United States. *BioScience* 57: 956-964.
- Edees, E.S. and A. Newton. 1988. *Brambles of the British Isles*. London: Ray Society.
- Flora of China Editorial Committee. 2003. *Flora of China*. Volume 9 (Pittosporaceae through Connaraceae) in eFloras (2008). Published on the Internet <http://www.efloras.org> [accessed 2019]. Missouri Botanical Garden, St. Louis, MO & Harvard University Herbaria, Cambridge, MA.
- _____. 2006. *Flora of China*. Vol. 22 (Poaceae) in eFloras (2008). Published on the Internet <http://www.efloras.org> [accessed 2019]. Missouri Botanical Garden, St. Louis, MO & Harvard University Herbaria, Cambridge, MA.
- Flora of North America Editorial Committee (FNA). 2010. *Flora of North America north of Mexico*. Volume 7, Magnoliophyta: Brassicaceae to Salicaceae. Oxford Univ. Press, New York, NY.
- _____. 2014. *Flora of North America north of Mexico*. Volume 9, Magnoliophyta: Picramniaceae to Rosaceae. Oxford Univ. Press, New York, NY.
- Green, A.F., T.S. Ramsey, and J. Ramsey. 2013. Polyploidy and invasion of English ivy (*Hedera* spp., Araliaceae) in North American forests. *Biological Invasions* 15: 2219-2241.
- Heffernan, K., E. Engle, and C. Richardson. 2014. Virginia invasive plant species list. Virginia Department of Conservation and Recreation, Division of Natural Heritage. Natural Heritage Technical Document 14-11. Richmond, Virginia.
- [Invasive Plants Species List - Virginia Department of Conservation ...](#)
- Belyaeva, Irina. 2009. Nomenclature of *Salix fragilis* L. and a new species, *S. euxina* (Salicaceae). *Taxon* 58(4): 1344-1348.
- McAllister, H. and A. Rutherford. 1990. *Hedera helix* L. and *H. hibernica* (Kirchner) Bean (Araliaceae) in the British Isles. *Watsonia* 18: 7-15.
- _____, and R.H. Simmons. In prep. Identification of *Hedera* (Araliaceae) in the mid-Atlantic and southeastern states.
- McAvoy, W.A. 2018. Non-native invasive plants of Delaware. Delaware Dept. of Natural Resources and Environmental Control.
- Nesom, G.L. 2009. Taxonomic overview of *Ligustrum* (Oleaceae) naturalized in North America north of Mexico. *Phytologia* 91: 467-482.
- Niemiera, A.X. 2010. Bradford callery pear (and other cultivars). Virginia Polytechnic Institute and State University, Blacksburg, Virginia. www.ext.vt.edu
- Pimentel D, R. Zuniga, and D. Morrison. 2005. Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics* 52: 273–288.
- Rickel, C. 2013. Field guide to identification of *Phyllostachys* invasive running bamboo. Institute of Invasive Bamboo Research.
- Simmons, R.H. 2015. Native vascular flora of the City of Alexandria, Virginia. City of Alexandria Department Recreation, Parks, and Cultural Activities, Alexandria, VA.
- <http://alexandriava.gov/22560>
- _____. 2010. Annals of the City of Alexandria herbarium: local range extensions in Orchidaceae. City of Alexandria Dept. Recreation, Parks, and Cultural Activities. Alexandria, Virginia.
- _____, M.T. Strong, and J.M. Parrish. 2008. Noteworthy Collections: Virginia. *Castanea* 73: 328-331.

- Steury, B.W. 2011. Additions to the vascular flora of the George Washington Memorial Parkway, Virginia, Maryland, and the District of Columbia. *Banisteria* 37: 3-20.
- Strong, M.T. and R.H. Simmons. 2002. Noteworthy Collections: Maryland. *Castanea* 67: 120-121.
- Swearingen, J., B. Slattery, K. Reshetiloff, and S. Zwicker. 2010. Plant invaders of mid-Atlantic natural areas. 4th ed. National Park Service and U.S. Fish and Wildlife Service. Washington, D.C. 168 pp.
- Urbatsch, L.E., J.F. Pruski, and K.M. Neubig. 2013. *Youngia thunbergiana* (Crepidinae, Cichorieae, Asteraceae), a species overlooked in the North American flora. *Castanea* 78: 330-337.
- USDA, NRCS. 2019. The PLANTS Database (<http://plants.usda.gov>). National Plant Data Team, Greensboro, NC 27401-4901 USA.
- Valcárcel, V. and P. Vargas. 2010. Quantitative morphology and species delimitation under the general lineage concept: optimization for *Hedera* (Araliaceae). *American Journal of Botany* 97(9): 1555-1573.
- Virginia Botanical Associates (VBA). 2019. Digital Atlas of the Virginia Flora (<http://www.vaplantatlas.org>). c/o Virginia Botanical Associates, Blacksburg, Virginia.
- Vitousek, P.M., C.M. D'Antonio, L.L. Loope, and R. Westbrooks. 1996. Biological invasions as global environmental change. *American Scientist* 84: 218-228.
- Weakley, A.S., J.C. Ludwig, and J.F. Townsend. 2012. Flora of Virginia. Bland Crowder, ed. Foundation of the Flora of Virginia Project Inc., Richmond. Fort Worth: Botanical Research Institute of Texas Press.
- Whittemore, A.T. 2004. Sawtooth oak (*Quercus acutissima*, Fagaceae) in North America. *Sida: Contributions to Botany* 21: 447-454.
- Zheng, H., Y. Wu, J. Ding, D. Binion, W. Fu, and R. Reardon. 2004. Invasive plants established in the United States that are found in Asia and their associated natural enemies. Vol. 1. 147 pp.
- _____. 2005. Invasive plants established in the United States that are found in Asia and their associated natural enemies. Vol. 2. 174 pp.

Prepared by:

Rod Simmons
Natural Resource Manager / Plant Ecologist
Natural Resources Division
Department of Recreation, Parks & Cultural Activities
City of Alexandria, Virginia
2900-A Business Center Drive
Alexandria, VA 22314
Rod.Simmons@alexandriava.gov

Citation:

Simmons, R.H. 2019. Non-native invasive plants of the City of Alexandria, Virginia. City of Alexandria Dept. Recreation, Parks, and Cultural Activities, Alexandria, Virginia.

Revised February 2019